



11) Publication number:

0 605 020 A1

EUROPEAN PATENT APPLICATION

21 Application number: 93200772.7

(51) Int. Cl.5: A23D 7/00

22 Date of filing: 17.03.93

(43) Date of publication of application: 06.07.94 Bulletin 94/27

Designated Contracting States:
AT BE CH DE DK ES FR GB GR IE IT LI NL PT
SE

Applicant: UNILEVER N.V.
P.O. Box 760
NL-3000 DK Rotterdam(NL)

BE CH DE DK ES FR GR IT LI NL PT SE AT

 Applicant: UNILEVER PLC Unilever House Blackfriars
 P.O. Box 68 London EC4P 4BQ(GB)

⊗ GB IE

Inventor: Schotel, Ronald Albert, UNILEVER RESEARCH LABORATORY VLAARDINGEN, Olivier van Noortlaan 120 NL-3133 AT Vlaardingen(NL)

Representative: Joppe, Hermina L. P. et al Unilever N.V. Patent Division P.O. Box 137 NL-3130 AC Vlaardingen (NL)

- 54 Spread containing oligofructoses.
- 67) A water-continuous spread comprising:
 - (A) a structuring amount of oligofructoses; and

EP 0 605 020 A1

The present invention is concerned with spread products, in particular with spread products of low fat content, and processes for their preparation.

Many attempts have been made to formulate low fat spread products. Amongst the various reasons why such products are desired is the wish to reduce the caloric content of the spread and other dietetic considerations.

A substance that has widely been applied as a fat extender is water. This use of water has, for example, led to the introduction of so-called halvarines. If relatively high levels of water are used, often thickening agents and/or gelling agents are used for avoiding adverse effects of the high water level.

For example EP 298 561 (Unilever N.V.) describes the preparation of edible plastic dispersions not having a continuous fat phase, said composition including at least two gelling agents forming two gelforming compositions.

EP 509 707 discloses low fat spreads containing a combination of water soluble starch, alginate and ion source. Spreads according to this document may optionally contain 1 to 15 % of a soluble vegetable fibre. An example of a suitable fibre for this purpose is INULIN which is for example marketed under the trade name FIBRULINE.

JP 267450/90 (Ajinimoto Co. Ltd.) discloses the use of polyfructan as a fat or oil substitute in food products. Preferred polyfructans are low caloric polysaccharides of the inulin type, which are mainly composed of beta-2,1-bonds).

Polysaccharides of the Inulin type are for example marketed under the trade name Raftiline™ and Fibruline™, these materials may advantageously be used in food products because they favourably contribute to the fibre content. A problem with the use of oligofructose materials is that when they are used in spreads, especially at high levels, sometimes they give rise to an undesired sweet off-taste. Also the appearance of spreads with relatively high levels of oligofructose materials is not always optimal.

It is the object of the present invention to formulate low fat spread products having a good flavour, texture and appearance. Surprisingly it has been found that these high quality, low fat products can be obtained if specific levels of oligofructose materials are used in combination with a narrowly defined level of mono- or disaccharides.

Accordingly the present invention relates to a water-continuous spread comprising:

- (A) a structuring amount of oligofructoses; and
- (B) from 0.1 to 15 wt% of added mono- or di-saccharides.

The improved taste of the product of the invention is especially surprising, because it appears that the addition of mono- or disaccharides to a spread containing structuring amounts of oligofructoses does not lead to an increase of the sweetness but rather to a decrease of the sweetness. Although applicants do by no means wish to be bound by this theory, it is believed that this taste-effect may be explained by the fact that the added mono- or di-saccharides have a masking effect, due to which the sweet taste of the oligofructose material is masked and the final product is perceived as less sweet as the product without the added mono-or di-saccharides.

For the purpose of the invention, preferably oligofructoses are used. Most preferred are materials containing one (terminal) sucrose unit, the remaining groups mainly being fructose units. Also the oligofructose is preferably of the beta-2,1 type, more specifically it is inulin which may be obtained from plants such as e.g. Compositae species and fructans obtained from various micro-organism as e.g. Aspergillus sydowii. Especially preferably inulin materials derived from Jerusalem artichoke of chicory are used. Preferably commercially available inulin materials like Raftiline or Fibruline are used. Other suitable oligofructoses are e.g. irisin and lycorisin.

The (weight) average degree of polymerisation of the oligofructose material is preferably from 6 to 100, more preferred 7 to 30, most preferred 8 to 25.

Compositions of the invention contain a structuring amount of oligofructoses. For the purpose of the invention, structuring amounts of oligofructoses are levels at which the oligofructose material is no longer soluble in the system, but is present as discrete particles which may provide structure to the product. The structuring amount of oligofructoses can suitably be determined by any suitable technique, for example by mixing all ingredients other than oligofructoses and then gradually adding oligofructoses while measuring the viscosity. At a certain level of oligofructoses a structure is formed by the oligofructoses. This may be evidenced e.g. by a change in melting temperature or an increase of viscosity. For the purpose of the invention any levels of oligofructoses above the level where a structure begins to form are referred to a structuring amounts.

The structuring amount of oligofructoses may depend on the type of oligofructoses used and the remaining ingredients of the product. For inulin type oligofructoses, in particular those derived from Jerusalem artichoke or chicory, the structuring amount is generally more than 15 wt%, for example 17 to 70

45

55

5

10

15

wt%, more preferred 20 to 50 wt%. For other oligofructoses similar structuring amounts apply.

Accordingly in a second embodiment of the invention there is provided a water-continuous spread comprising:

- (A) more than 15 wt% of oligofructoses; and
- (B) from 0.1 to 15 wt% of added mono- or disaccharides.

Preferably the particle size of the oligofructose agglomerates in the spread is in the order of magnitude of 0.5 - 20, preferably 1 - 5 micrometer which can be effected by suitable processing. The size of the primary oligofructose particles preferably is in the order of magnitude of 50 to 500 nm.

Compositions of the invention contain from 0.1 to 15 wt% of added mono- or disaccharides, more preferably 0.5 to 10 wt%, most preferred 0.7 to 5 wt%. For the purpose of the invention added mono- or disaccharides are those mono- or di-saccharides which are added on top of the mono- or di-saccharides which may be present in the oligofructose material.

Suitable mono-saccharides are for example Ribose, Arabinose, Xylose, Lyxose, Sorbitol, Glucose, Mannose and Fructose. Suitable di-saccharides are for example Isomaltose, Lactose, Maltose, Melibiose and Sucrose. Preferably the added mono- or di-saccharides are other mono- or di-saccharides than those which may be present in the oligofructose materials (mainly glucose, fructose and sucrose). Also preferably di-saccharides are used, most preferred is the use of Lactose.

Lactose can suitably derived from a dairy source, for example from whole milk, semi-skimmed milk, skimmed milk, (cultured) buttermilk, butter milk powder, skimmed milk powder, (Greek) yoghurt, quark, (low fat) fromage frais, (low fat) cottage cheese, butter, (Philadelphia) cream cheese, (double) cream, Ricotta cheese etc. In a preferred embodiment of the invention the lactose is derived from dairy cream or cream cheese.

In another preferred embodiment of the invention the di-saccharide is derived from non-dairy creams. Generally non-dairy creams contain lactose as disaccharide material.

Compositions of the invention may optionally comprise from 0.1 to 7 wt% of fat, more preferred the level of fat is more than 0.5 wt% and less than 4 wt%, most preferred are fat levels from 1 to 3 wt%.

Throughout this specification the terms oil and fat are used interchangeably. They are meant to include triglycerides of natural or synthetic origin such as soybean oil, sunflower oil, palm oil, fish oil, rapeseed oil, coconut oil, and hydrogenated, fractionated and/or interesterified triglyceride mixtures as well as edible substances that are physically similar to triglycerides such as waxes, e.g. jojoba oil and poly fatty acid esters of mono- or di-saccharides, and that can be used as replacement for or in admixture with triglycerides.

Preferably the fat is present in the form of small fat globules or crystals, having a volume weighted mean diameter of less than 20 μ m, more preferred from 0.1 to 5 μ m, most preferred from 0.5 to 2 μ m. The presence of these small fat particles in low fat spreads of the invention has the distinct advantage of providing an improved appearance to the product.

In a very preferred embodiment of the invention at least part of the fat is dairy fat. Preferably at least 10 wt% based on the total weight of the fat is dairy fat, more preferred more than 50 wt%, most preferred more than 90 wt% or even 100 wt%. The dairy fat can be derived from any dairy source as mentioned above. Especially preferably the dairy fat is derived from dairy cream or cream cheese.

In another preferred embodiment of the invention the fat is derived from non-dairy creams.

The use of fat sources wherein the ratio of fat to protein is high, for example butterfat, cream cheese and dairy cream or non-dairy cream, is preferred. These materials provide an excellent flavour to the product. Preferably fat sources are used, wherein the weight ratio of fat to protein is more than 5:1, more preferred more than 10:1, most preferred from 50:1 to 1000:1.

For optimising the flavour of the spreads of the invention it is especially preferred that the di-saccharide material or the dairy fat is incorporated in the product by using at least 5 wt% of a dairy ingredient in the spread. The optimum level of dairy ingredients will be dependent on the type and saccharide or fat level of the dairy product. Also combinations of dairy products may be used.

If whole milk, semi skimmed milk, skimmed milk or combinations thereof are used, the total level thereof is preferably from 40 to 85 wt% of the composition, more preferred 50-80 wt%, most preferred 55-80 wt%.

If yoghurt, quark, cottage cheese, butter fat, cream, cream cheese or fromage frais or a combination thereto is used, the total level is preferably from 2-40 wt%, more preferred 5-30 wt%. Under some circumstances it may be advantageous to use a mixture of milk and these ingredients, for example in weight ratios between 20 : 1 and 2 : 1, the total level of yoghurt/quark/cottage cheese/fromage frais/cream/cream cheese and milk being from 60-85 wt%.

If non-dairy cream is used, the preferred level is from 2 to 40 wt%, more preferred 5 to 30 wt%.



Especially preferred is the combined use of cream (dairy or non-dairy) with butter fat, for example in a weight ratio of 20 : 1 to 1 : 1.

Preferably spreads of the invention are plastic in the sense that they can be spread onto bread without tearing the bread. Generally plastic spreads will have a stress strain relation with a maximum stress occurring at strain of 0.001-0.5 (preferably 0.001 to 0.3), the maximum stress at this strain being 0.01-100 kPa and with a ratio of plastic stress and the maximum stress of 0.1 to 1. A suitable method for determining these values is given in EP 298 561.

In addition to the above mentioned ingredients, spreads of the invention may comprise a number of optional ingredients such as flavouring, salt, preservatives, acidifiers, vitamins, colouring materials etc.

Preferably the level of flavouring materials (other than those which are incorporated through the dairy ingredients) is less than 0.5 wt%, for example 0.01 to 2 wt%. Preferably the level of salt (sodium chloride) is from 0-4 wt%, more preferred 0.1 to 3 wt%, most preferred 0.5 to 1.2 wt%. Preservatives are preferably incorporated at a level of 0-4 wt%, more preferred 0.01 to 1 wt%, most preferred 0.05 to 0.3 wt%. Especially preferred is the use of potassium sorbate. A preferred colouring material is carotene; preferred levels of colouring material are from 0 - 1 wt%, more preferred 0.01 to 0.2 wt%. Acidifiers may be incorporated to bring the pH of the product to the desired level, preferably the pH of the product is from 3 to 10, more preferred 3.5 to 7. A suitable acidifier is for example lactic acid or citric acid.

Another optional ingredient which may be present in compositions of the invention are proteins. Preferably the protein level in spreads of the invention is from 0.5 to 15 wt%, more preferred, 2 to 6 wt%, most preferred 2.5 to 4 %. In an especially preferred embodiment of the invention the protein are partially or wholly obtained from dairy sources. In another preferred embodiment of the invention the protein is wholly or partially a vegetable protein, especially soy bean protein. For example if mixtures of these ingredients are used suitable weight ratios of dairy protein to vegetable protein may for example be from 10: 1 to 1:10.

The spread may further containing small particles such as herbs and vegetables. The total level thereof will generally be less than 10 wt%.

Spreads of the invention will generally comprise fairly high levels of water, say from 50 to 84.9 wt% of the composition, more preferred 60 to 80 wt%, most preferred 65 to 75 wt%. Water may be incorporated as such, or as part of the other ingredients such as milk etc. Spreads of the invention are water-continuous in the sense that they comprise a continuous aqueous phase. Water-continuous spreads can readily be distinguished from fat-continuous spreads by virtue of conductivity measurements.

Spreads of the invention can suitably be used as bread spreads to replace e.g. margarine or halvarine, but they can also suitable be used as flavoured spread, for example cheese spreads, meat spreads, nut spreads, sweet spreads or vegetable spreads.

Spreads of the invention may be prepared by any suitable method for the preparation of water-continuous spread products. A preferred method however involves the mixing of the ingredients, optionally followed by pasteurisation and homogenisation, followed by cooling in one or more scraped surface heat exchangers (A-units) to a temperature of 0 to 10 °C, optionally followed by processing through one or more C-units. Pasteurisation is especially preferred for extending the keepability of the product. Homogenisation is preferred for effecting the formation of small fat globules to enhance the appearance of the spread. After preparation the spreads are generally packed in wrappers or containers, generally tubs with a content of 2-1000 g will be used, especially 5, 10, 15, 250 or 500 g. Preferably the packed product is stored at refrigerator temperatures.

45

25

50

EXAMPLES

The following compositions were made:

10	

15

5

Ingredient wt%	Α	В	С	D
Inulin	33	33	33	33
Butter fat	-	1	-	
Non-dairy cream	12	12	_	
Dairy cream	-	-	12	-
Potassium sorbate	0.12	0.12	0.12	0.12
Salt	0.8	0.8	0.8	0.8
Flavouring	150 ppm	150 ppm	150 ppm	150 ppm
Lactic acid	to pH 4.8	to pH 4.8	to pH 4.8	to pH 4.8
Water	balance	balance	balance	balance

The inulin was Raftiline LS ex Tiense Suiker raffinaderij The non-dairy cream was Lactofil classic containing 26.7 wt% fat, 2 wt% protein and 19.9 wt% lactose. The dairy cream contained 45 wt% fat.

The product was prepared as follows: the cream and water were mixed at 60 °C, cooled to 45 °C whereafter the pH was adjusted and the remaining ingredients were added.

Products A-C (according to the invention) had an improved taste and appearance when compared to product D (comparison).

Claims

- 1. A water-continuous spread comprising:
 - (A) a structuring amount of oligofructoses; and
 - (B) from 0.1 to 15 wt% of mono- or di-saccharides.
- 2. A water-continuous spread comprising:
 - (A) more than 15 wt% of oligofructoses; and
 - (B) from 0.1 to 15 wt% of added mono- or disaccharides.
- 3. A spread according to claim 1 or 2, wherein the oligofructose is an inulin, preferably derived from Jerusalem artichoke or chicory.
 - 4. A spread according to claim 1 or 3, wherein the level of oligofructoses is more than 15 wt%, more preferred 17 to 70 wt%, most preferred 20 to 50 wt%.
- 5. A spread according to one or more of the preceding claims, wherein the saccharide is a di-saccharide, preferably lactose.
 - 6. A spread according to one or more of the preceding claims comprising from 0.1 to 7 wt% of fat.
- 7. A spread according to claim 4 or 5, wherein the di-saccharide and/or the fat is derived from a dairy source, preferably dairy cream or from non-dairy cream.

50



EUROPEAN SEARCH REPORT

Application Number

EP 93 20 0772 Page 1

Category	Citation of document with in of relevant par	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. CL.5)	
x	US-A-5 169 671 (TSU* column 2, line 3 claims 1,3,5,13,16, table 24 *	1,4,6,7	A23D7/00	
x	DATABASE WPI Week 9305, Derwent Publications Ltd., London, GB; AN 93-039127 & JP-A-4 356 169 (AJINOMOTO KK) 9 December 1992 * abstract *			
X	PATENT ABSTRACTS OF vol. 16, no. 106 (C & JP-A-32 80 857 (December 1991 * abstract *	JAPAN -919)16 March 1992 AJINOMOTO CO INC) 11	1	
E .	WO-A-9 306 744 (RAF S.A.) * claims 1-11,18-20	1-4	TECHNICAL FIELDS SEARCHED (Int. Cl.5)	
D, X	DATABASE WPI Week 9238, Derwent Publications Ltd., London, GB; AN 92-310708 & JP-A-4 210 578 (AJINOMOTO KK) 31 July 1992			A23D A23L A23C
א,ם	EP-A-0 509 707 (PET * page 3, line 3 - 1,4,5,9,16,20; exam	RELLA LIMITED) line 7; claims ples 1-5,8,9,11-14 *	1,3,5-7	
A	EP-A-0 388 572 (SER * claims 2,13 *	ES LABORATORIES, INC.) -/	1	
	The present search report has b			Remine
	Place of search THE HAGUE	Data of complation of the search 15 JULY 1993		DEKEIREL M.J.
X:pa Y:pa do	CATEGORY OF CITED DOCUME rticularly relevant if taken alone rticularly relevant if combined with an current of the same category chrological background	NTS T: theory or princi E: earlier patent de after the filing c other D: document cited L: document cited	ocument, but put date in the application for other reasons	blished on, or



EUROPEAN SEARCH REPORT

Application Number

EP 93 20 0772 Page 2

	P.O.O.I.			Page 2
	DOCUMENTS CONS	IDERED TO BE RELEV	ANT	
Category	Citation of document with of relevant p	indication, where appropriate,	Relevant to claim	CLASSIFICATION OF 1 APPLICATION (Int. CL.
A	PATENT ABSTRACTS OF		1	A PERCATION (IRI. CL.
		NIPPON OIL & FATS CO		
				TECHNICAL FIELDS SEARCHED (Int. CL5)
!				
	he present search report has been			
	HAGUE	Date of completion of the search 15 JULY 1993	DE	KEIREL M.J.
X : particul Y : particul docume A : technoli	EGORY OF CITED DOCUMENTS ariy relevant if taken sione ariy relevant if combined with anothe nt of the same category spical background	E: serier patent d after the filing D: document cited L: document cited	in the application for other reasons	of an, or
∪ : 1001. - 17 1	tion disclosure diste document	A : member of the	same patent family, o	Presponding

